

Subleading-twist effects in single-spin asymmetries in semi-inclusive deep-inelastic scattering on a longitudinally polarized hydrogen target

A. Airapetian,¹⁸ N. Akopov,³⁰ Z. Akopov,³⁰ M. Amarian,^{8,30} A. Andrus,¹⁶ E.C. Aschenauer,⁸ W. Augustyniak,²⁹ R. Avakian,³⁰ A. Avetissian,³⁰ E. Avetissian,¹² A. Bacchetta,²³ P. Bailey,¹⁶ D. Balin,²¹ M. Beckmann,⁷ S. Belostotski,²¹ N. Bianchi,¹² H.P. Blok,^{20,28} H. Böttcher,⁸ A. Borissov,¹⁵ A. Borysenko,¹² M. Bouwhuis,¹⁶ A. Brüll,¹⁷ V. Bryzgalov,²² M. Capiluppi,¹¹ G.P. Capitani,¹² T. Chen,⁴ G. Ciullo,¹¹ M. Contalbrigo,¹¹ P.F. Dalpiaz,¹¹ W. Deconinck,¹⁸ R. De Leo,³ M. Demey,²⁰ L. De Nardo,¹ E. De Sanctis,¹² E. Devitsin,¹⁹ M. Diefenthaler,¹⁰ P. Di Nezza,¹² J. Dreschler,²⁰ M. Düren,¹⁴ M. Ehrenfried,¹⁰ A. Elalaoui-Moulay,² G. Elbakian,³⁰ F. Ellinghaus,⁶ U. Elschenbroich,¹³ R. Fabbri,²⁰ A. Fantoni,¹² L. Felawka,²⁶ S. Frullani,²⁴ A. Funel,¹² G. Gapienko,²² V. Gapienko,²² F. Garibaldi,²⁴ K. Garrow,²⁶ G. Gavrillov,^{7,26} V. Gharibyan,³⁰ O. Grebeniouk,²¹ I.M. Gregor,⁸ C. Hadjidakis,¹² K. Hafidi,² M. Hartig,¹⁴ D. Hasch,¹² W.H.A. Hesselink,^{20,28} A. Hillenbrand,¹⁰ M. Hoek,¹⁴ Y. Holler,⁷ B. Hommez,¹³ I. Hristova,⁸ G. Iarygin,⁹ A. Ivanilov,²² A. Izotov,²¹ H.E. Jackson,² A. Jgoun,²¹ R. Kaiser,¹⁵ T. Keri,¹⁴ E. Kinney,⁶ A. Kisselev,^{6,21} T. Kobayashi,²⁷ M. Kopytin,⁸ V. Korotkov,²² V. Kozlov,¹⁹ B. Krauss,¹⁰ V.G. Krivokhijine,⁹ L. Lagamba,³ L. Lapikás,²⁰ A. Laziev,^{20,28} P. Lenisa,¹¹ P. Liebing,⁸ L.A. Linden-Levy,¹⁶ W. Lorenzon,¹⁸ H. Lu,⁵ J. Lu,²⁶ S. Lu,¹⁴ B.-Q. Ma,⁴ B. Maiheu,¹³ N.C.R. Makins,¹⁶ Y. Mao,⁴ B. Marianski,²⁹ H. Marukyan,³⁰ F. Masoli,¹¹ V. Mexner,²⁰ N. Meyners,⁷ T. Michler,¹⁰ O. Mikloukhov,²¹ C.A. Miller,^{1,26} Y. Miyachi,²⁷ V. Muccifora,¹² M. Murray,¹⁵ A. Nagaitsev,⁹ E. Nappi,³ Y. Naryshkin,²¹ M. Negodaev,⁸ W.-D. Nowak,⁸ K. Oganessyan,^{7,12} H. Ohsuga,²⁷ A. Osborne,¹⁵ N. Pickert,¹⁰ D.H. Potterveld,² M. Raithel,¹⁰ D. Reggiani,¹⁰ P.E. Reimer,² A. Reischl,²⁰ A.R. Reolon,¹² C. Riedl,¹⁰ K. Rith,¹⁰ G. Rosner,¹⁵ A. Rostomyan,³⁰ L. Rubacek,¹⁴ J. Rubin,¹⁶ D. Ryckbosch,¹³ Y. Salomatin,²² I. Sanjiev,^{2,21} I. Savin,⁹ A. Schäfer,²³ G. Schnell,^{8,27} K.P. Schüller,⁷ J. Seele,⁶ R. Seidl,¹⁰ B. Seitz,¹⁴ C. Shearer,¹⁵ T.-A. Shibata,²⁷ V. Shutov,⁹ K. Sinram,⁷ W. Sommer,¹⁴ M. Stancari,¹¹ M. Statera,¹¹ E. Steffens,¹⁰ J.J.M. Steijger,²⁰ H. Stenzel,¹⁴ J. Stewart,⁸ F. Stinzing,¹⁰ P. Tait,¹⁰ H. Tanaka,²⁷ S. Taroian,³⁰ B. Tchuiko,²² A. Terkulov,¹⁹ A. Trzcinski,²⁹ M. Tytgat,¹³ A. Vandenbroucke,¹³ P.B. van der Nat,²⁰ G. van der Steenhoven,²⁰ Y. van Haarlem,¹³ V. Vikhrov,²¹ M.G. Vinther,¹ C. Vogel,¹⁰ J. Volmer,⁸ S. Wang,⁴ J. Wendland,^{25,26} Y. Ye,⁵ Z. Ye,⁷ S. Yen,²⁶ B. Zihlmann,¹³ and P. Zupranski²⁹

(The HERMES Collaboration)

¹*Department of Physics, University of Alberta, Edmonton, Alberta T6G 2J1, Canada*

²*Physics Division, Argonne National Laboratory, Argonne, Illinois 60439-4843, USA*

³*Istituto Nazionale di Fisica Nucleare, Sezione di Bari, 70124 Bari, Italy*

⁴*School of Physics, Peking University, Beijing 100871, China*

⁵*Department of Modern Physics, University of Science and Technology of China, Hefei, Anhui 230026, China*

⁶*Department of Physics, University of Colorado, Boulder, Colorado 80309-0390, USA*

⁷*DESY, 22603 Hamburg, Germany*

⁸*DESY, 15738 Zeuthen, Germany*

⁹*Joint Institute for Nuclear Research, 141980 Dubna, Russia*

¹⁰*Physikalisches Institut, Universität Erlangen-Nürnberg, 91058 Erlangen, Germany*

¹¹*Istituto Nazionale di Fisica Nucleare, Sezione di Ferrara and*

Dipartimento di Fisica, Università di Ferrara, 44100 Ferrara, Italy

¹²*Istituto Nazionale di Fisica Nucleare, Laboratori Nazionali di Frascati, 00044 Frascati, Italy*

¹³*Department of Subatomic and Radiation Physics, University of Gent, 9000 Gent, Belgium*

¹⁴*Physikalisches Institut, Universität Gießen, 35392 Gießen, Germany*

¹⁵*Department of Physics and Astronomy, University of Glasgow, Glasgow G12 8QQ, United Kingdom*

¹⁶*Department of Physics, University of Illinois, Urbana, Illinois 61801-3080, USA*

¹⁷*Laboratory for Nuclear Science, Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA*

¹⁸*Randall Laboratory of Physics, University of Michigan, Ann Arbor, Michigan 48109-1040, USA*

¹⁹*Lebedev Physical Institute, 117924 Moscow, Russia*

²⁰*Nationaal Instituut voor Kernfysica en Hoge-Energiefysica (NIKHEF), 1009 DB Amsterdam, The Netherlands*

²¹*Petersburg Nuclear Physics Institute, St. Petersburg, Gatchina, 188350 Russia*

²²*Institute for High Energy Physics, Protvino, Moscow region, 142281 Russia*

²³*Institut für Theoretische Physik, Universität Regensburg, 93040 Regensburg, Germany*

²⁴*Istituto Nazionale di Fisica Nucleare, Sezione Roma 1, Gruppo Sanità and Physics Laboratory, Istituto Superiore di Sanità, 00161 Roma, Italy*

²⁵*Department of Physics, Simon Fraser University, Burnaby, British Columbia V5A 1S6, Canada*

²⁶*TRIUMF, Vancouver, British Columbia V6T 2A3, Canada*

²⁷*Department of Physics, Tokyo Institute of Technology, Tokyo 152, Japan*

²⁸*Department of Physics and Astronomy, Vrije Universiteit, 1081 HV Amsterdam, The Netherlands*

²⁹Andrzej Soltan *Institute for Nuclear Studies, 00-689 Warsaw, Poland*
³⁰Yerevan Physics Institute, 375036 Yerevan, Armenia
 (Dated: February 4, 2008)

Single-spin asymmetries in the semi-inclusive production of charged pions in deep-inelastic scattering from transversely and longitudinally polarized proton targets are combined to evaluate the subleading-twist contribution to the longitudinal case. This contribution is significantly positive for π^+ mesons and dominates the asymmetries on a longitudinally polarized target previously measured by HERMES. The subleading-twist contribution for π^- mesons is found to be small.

PACS numbers: 13.60.-r, 13.88.+e, 14.20.Dh, 14.65.-q

Single-spin asymmetries in the distribution of lepton-produced hadrons in the azimuthal angle around the virtual photon direction are a valuable tool for the exploration of transverse spin and momentum degrees of freedom in nucleon structure. Whereas two out of the three fundamental quark distributions, the unpolarized quark density and the helicity density, can be accessed in inclusive measurements, this is not true for the remaining and so far unmeasured transversity distribution function [1, 2, 3]. Since transversity is chiral-odd and since hard interactions conserve chirality, it can only be probed by a process involving some additional chiral-odd object. Single-spin asymmetries in semi-inclusive deep-inelastic scattering (SIDIS), e.g., involving the chiral-odd Collins fragmentation function [4], could be the required quark “polarimeter” to access transversity. This has been a main motivation to look for azimuthal single-spin asymmetries. Such asymmetries have been observed in SIDIS with unpolarized beams and with targets polarized both longitudinally and transversely with respect to the beam direction [5, 6, 7, 8, 9]. Asymmetries have also been observed with polarized beams and unpolarized nucleons [10, 11]. The asymmetry for a transversely polarized target can be interpreted in terms of the transversity distribution function, convoluted with the Collins fragmentation function, as well as in terms of the Sivers [12] function, which appears with the ordinary unpolarized fragmentation function. In the case of targets that are polarized longitudinally with respect to the incoming beam direction, the interpretation is more complex. In fact, the asymmetry for a target polarized along the virtual photon direction contains various contributions from subleading-twist quark distribution and fragmentation functions. These contributions have – for dynamics reasons – an additional $1/Q$ suppression compared to the ordinary $1/Q^4$ suppression of the Mott cross section. When the polarization is along the beam direction, the small but non-vanishing component of the nuclear spin transverse to the photon direction, although $1/Q$ suppressed for kinematical reasons, also contributes to the measured asymmetry. This feature has been exploited in several estimates [13, 14, 15, 16, 17, 18, 19] for the hitherto unknown transversity distribution and Collins fragmentation functions. However, some or all of the above-mentioned subleading-twist terms have been




FIG. 1: The definitions of the azimuthal angle ϕ of the hadron production plane, relative to the plane containing the momentum \mathbf{l} (\mathbf{l}') of the incident (scattered) lepton, the polar angle θ_{γ^*} between the virtual photon and the incoming lepton directions, and of the transverse component \mathbf{S}_{\perp} of the target spin \mathbf{S} with respect to the photon direction $\mathbf{q} \equiv \mathbf{l} - \mathbf{l}'$.

neglected in all these estimates.

In this Letter the recently measured asymmetries on a transversely polarized hydrogen target [8] are used to eliminate the contribution due to the transverse spin component from the measured asymmetries on a longitudinally polarized hydrogen target [5], thereby allowing, for the first time, the extraction of the purely subleading-twist contribution. Knowledge of this subleading contribution is essential to any extraction of information on the transversity distribution or Collins fragmentation function from data with longitudinal target polarization.

Whenever the target is polarized with respect to the incoming beam direction the measured asymmetries contain contributions from both the transverse and longitudinal polarization components with respect to the virtual photon direction. Throughout this paper asymmetries and their azimuthal moments will carry one of the following superscripts for distinction: “q” when the reference axis is the photon direction and “l” when it is the lepton beam. They will be called photon-axis or lepton-axis asymmetries/moments, respectively. Asymmetries and moments will also carry two-letter subscripts denoting the polarization of beam and target. For the definition of azimuthal angles, asymmetries, and azimuthal moments thereof the *Trento Conventions* [20] will be used.

The size of the component of the nucleon spin vector that is transverse to the virtual photon direction depends on θ_{γ^*} , the polar angle between the incoming beam direction and the virtual photon direction (see Fig. 1). Hence it strongly depends on the event kinematics. At HERMES kinematics $\sin\theta_{\gamma^*}$ can be as large as 15%. In the configuration shown in Fig. 1, where in the lab frame the target spin vector is opposite to the incoming beam direction, the transverse spin component lies in the lepton scattering plane. Therefore ϕ_S , the azimuthal angle of the target spin relative to the scattering plane, is equal to π .

The azimuthal moments of the distribution of hadrons around the virtual-photon direction can be separated into contributions from the longitudinal and transverse com-

ponents of the target polarization with respect to the virtual photon direction. For the case of the longitudinal lepton-axis moment $\langle \sin\phi \rangle_{UL}^1$, the contributions from the transverse component are the Sivers and Collins moments $\langle \sin(\phi - \phi_S) \rangle_{UT}^q$ and $\langle \sin(\phi + \phi_S) \rangle_{UT}^q$. Since in this case $\phi_S = \pi$, both moments contribute to the $\sin\phi$ Fourier component of the longitudinal lepton-axis asymmetry with a minus sign. In case of a transversely polarized target contributions arise from the dominating transverse and from the small but non-vanishing longitudinal component. Both the measured lepton-axis moments $\langle \sin(\phi + \phi_S) \rangle_{UT}^1$ and $\langle \sin(\phi - \phi_S) \rangle_{UT}^1$ contain contributions from $\langle \sin\phi \rangle_{UL}^q$. The lepton-axis moments are related to the photon-axis moments via the equation

$$\begin{pmatrix} \langle \sin\phi \rangle_{UL}^1 \\ \langle \sin(\phi - \phi_S) \rangle_{UT}^1 \\ \langle \sin(\phi + \phi_S) \rangle_{UT}^1 \end{pmatrix} = \begin{pmatrix} \cos\theta_{\gamma^*} & -\sin\theta_{\gamma^*} & -\sin\theta_{\gamma^*} \\ \frac{1}{2}\sin\theta_{\gamma^*} & \cos\theta_{\gamma^*} & 0 \\ \frac{1}{2}\sin\theta_{\gamma^*} & 0 & \cos\theta_{\gamma^*} \end{pmatrix} \begin{pmatrix} \langle \sin\phi \rangle_{UL}^q \\ \langle \sin(\phi - \phi_S) \rangle_{UT}^q \\ \langle \sin(\phi + \phi_S) \rangle_{UT}^q \end{pmatrix}, \quad (1)$$

which is valid up to corrections of order $\sin^2\theta_{\gamma^*}$ [21].

The complete analysis up to subleading-twist and leading order in α_s of longitudinal single-spin asymmetries in semi-inclusive DIS was presented in Ref. [22], completing

previous work of Refs. [23, 24]. Neglecting quark mass effects, the first term of the photon-axis moments on the right-hand-side of Eq. (1) is

$$\langle \sin\phi \rangle_{UL}^q = -\frac{(2-y)\sqrt{1-y}}{1-y+\frac{y^2}{2}} \frac{M}{Q} \frac{\mathcal{I}\left[\frac{\hat{\mathbf{P}}_{h\perp} \cdot \mathbf{k}_T}{M_h} \left(\frac{M_h}{zM} g_{1L} G^\perp + x h_L H_1^\perp\right) + \frac{\hat{\mathbf{P}}_{h\perp} \cdot \mathbf{p}_T}{M} \left(\frac{M_h}{zM} h_{1L}^\perp \tilde{H} - x f_L^\perp D_1\right)\right]}{f_1 D_1}. \quad (2)$$

The shorthand notation $\mathcal{I}[\mathcal{W}fD]$ is used here for the convolution integral appearing in the SIDIS cross section when quark transverse momenta are included, i.e.,

$$\mathcal{I}[\mathcal{W}fD] \equiv \int d^2\mathbf{P}_{h\perp} d^2\mathbf{p}_T d^2\mathbf{k}_T \delta^{(2)}\left(\mathbf{p}_T - \frac{\mathbf{P}_{h\perp}}{z} - \mathbf{k}_T\right) \times [\mathcal{W}f(x, p_T^2) D(z, z^2 k_T^2)], \quad (3)$$

where $\mathbf{P}_{h\perp}$ is the transverse momentum of the detected hadron, \mathbf{p}_T (\mathbf{k}_T) is the intrinsic quark transverse momentum in the generic distribution function f (fragmentation function D), and \mathcal{W} is a weight that depends on the involved distribution and fragmentation functions. The masses m_q , M , and M_h are the quark, nucleon and hadron masses and x , y , and z are the usual semi-inclusive DIS Lorentz invariants. The quark charge squared weighted sum over the various (anti)quark fla-

vors and the dependence on x (z), p_T^2 (k_T^2), and Q^2 of the distribution (fragmentation) functions have been omitted in Eq. (2).

Since the extraction of the subleading-twist term $\langle \sin\phi \rangle_{UL}^q$ is the main result of this work, its components are described briefly. The asymmetry arises from the interference of the scattering amplitudes of a longitudinal and transverse photon. This leads to the specific dependence of the numerator on the variable y . All terms in the numerator of Eq. (2) involve either combinations of subleading-twist distribution functions (h_L , f_L^\perp) with leading-twist fragmentation functions or of leading-twist distribution functions in conjunction with subleading-twist fragmentation functions (G^\perp , \tilde{H}). One should note that it is not possible to give a simple probabilistic interpretation to the subleading-twist functions. The terms

containing h_L and h_{1L}^\perp have been studied in some detail in Refs. [14, 15], making use of Wandzura-Wilczek approximations, and of Lorentz covariance relations. (The latter have been proven to be not rigorous in Ref. [25].) Note that the function h_{1L}^\perp appears also in the $\sin 2\phi$ Fourier component of the longitudinal single-spin asymmetry [23]. Recent preliminary results of CLAS support a non-vanishing $\sin 2\phi$ moment [26]. However, in measurements at HERMES which were in a different kinematic region than the ones at CLAS, it was found to be consistent with zero [5]. The term with the helicity distribution g_{1L} contains the fragmentation function G^\perp , which is at present unknown. A similar term appears also in the longitudinal beam-helicity asymmetry, A_{LU} [22]. The latter has been found to be non-zero both at HERMES [11] and at CLAS [10]. Finally, the last term of the numerator contains the function f_L^\perp . Similar to the Siverson function f_{1T}^\perp it is odd under time reversal (T-odd) and for this reason has been neglected in virtually all theoretical treatments of the measured lepton-axis asymmetries on a longitudinally polarized target. Recently, it has been recognized that such T-odd distribution functions can arise through initial or final state interactions [27, 28, 29, 30]. A calculation of the function f_L^\perp has been performed in Ref. [31] in the context of a simple diquark spectator model. It should be noted that so far factorization has been proven only for leading-twist observables in semi-inclusive deep-inelastic scattering with hadrons in the current fragmentation region detected at low transverse momentum [32, 33]. A factorization proof for subleading-twist observables is still open. At the moment no firm experimental information about any of the subleading-twist terms in Eq. (2) exist.

The extraction of $\langle \sin \phi \rangle_{UL}^q$ reported here gives a first indication about the size of such subleading-twist effects in azimuthal target-spin asymmetries. This is especially important when measuring leading-twist asymmetries of the same order of magnitude where the question arises whether or not subleading-twist contributions can be neglected.

The other two terms on the right-hand-side of Eq. (1) read

$$\langle \sin(\phi - \phi_S) \rangle_{UT}^q = - \frac{\mathcal{I} \left[\frac{\hat{\mathbf{p}}_{h_1} \cdot \mathbf{p}_T}{2M} f_{1T}^\perp D_1 \right]}{f_1 D_1}, \quad (4)$$

$$\langle \sin(\phi + \phi_S) \rangle_{UT}^q = - \frac{1-y}{1-y+\frac{y^2}{2}} \frac{\mathcal{I} \left[\frac{\hat{\mathbf{p}}_{h_1} \cdot \mathbf{k}_T}{2M_h} h_1 H_1^\perp \right]}{f_1 D_1} \quad (5)$$

which are the leading-twist Siverson and Collins moments, involving either the Siverson distribution function, or the transversity distribution h_1 in conjunction with the Collins fragmentation function H_1^\perp .

Measurements of azimuthal single-spin asymmetries on a transversely polarized target can be used to eliminate the contribution of these leading-twist mo-

ments (4) and (5) to the longitudinal lepton-axis moment $\langle \sin \phi \rangle_{UL}^1$. Hence the subleading-twist terms (Eq. (2)) can be isolated. At HERMES kinematics the deviation from unity of $\cos \theta_{\gamma^*}$ can be neglected. The subleading-twist contribution $\langle \sin \phi \rangle_{UL}^q$ then reads

$$\begin{aligned} \langle \sin \phi \rangle_{UL}^q &= \langle \sin \phi \rangle_{UL}^1 + \sin \theta_{\gamma^*} \langle \sin(\phi + \phi_S) \rangle_{UT}^1 \\ &\quad + \sin \theta_{\gamma^*} \langle \sin(\phi - \phi_S) \rangle_{UT}^1. \end{aligned} \quad (6)$$

Here $\sin \theta_{\gamma^*}$ is evaluated from the lepton kinematics as $2xMQ^{-1} \sqrt{1-y-y^2x^2M^2/Q^2} / \sqrt{1+4x^2M^2/Q^2}$.

For the extraction of the subleading-twist contribution $\langle \sin \phi \rangle_{UL}^q$ according to Eq. (6) the lepton-axis asymmetries on a longitudinally polarized hydrogen target [5] were reanalyzed to have the same binning in x and z as in the measurement on a transversely polarized hydrogen target [8]. Furthermore, the $\sin \phi$ modulation of the semi-inclusive cross section has been extracted in a fit of the normalized-yield asymmetry¹

$$A_{UL}^1(\phi) = \frac{1}{|P_L|} \frac{N^+(\phi) - N^-(\phi)}{N^+(\phi) + N^-(\phi)}, \quad (7)$$

where P_L is the longitudinal target polarization and \rightarrow / \leftarrow denotes a target polarized antiparallel/parallel to the incoming beam direction. The same requirements on the lepton kinematics were used as in the analysis of the transverse target data, i.e., $W^2 > 10 \text{ GeV}^2$, $0.023 < x < 0.4$, $0.1 < y < 0.85$ and $Q^2 > 1 \text{ GeV}^2$, where W is the invariant mass of the initial photon-nucleon system. Coincident hadrons were accepted only if $0.2 < z < 0.7$ and $\theta_{\gamma^*h} > 0.02 \text{ rad}$, where θ_{γ^*h} is the angle between the directions of the virtual photon and the hadron. The latter requirement was imposed to avoid a region where the azimuthal angle ϕ is poorly reconstructed due to detector smearing. Pions were identified in the momentum range $4 \text{ GeV} < P_\pi < 13.8 \text{ GeV}$ using either a threshold Cherenkov counter for the longitudinally polarized data set or a Ring Imaging Cherenkov counter for the transversely polarized target data. The lepton-axis moments from the transversely polarized target data set were extracted in the fit

$$\begin{aligned} A_{UT}^1(\phi, \phi_S) &= 2 \langle \sin(\phi + \phi_S) \rangle_{UT}^1 \sin(\phi + \phi_S) \\ &\quad + 2 \langle \sin(\phi - \phi_S) \rangle_{UT}^1 \sin(\phi - \phi_S) \end{aligned} \quad (8)$$

of the transverse asymmetry in Eq. (1) of Ref. [8].²

¹ This is in contrast to the previous publication on longitudinal single-spin asymmetries [5] where a weighting method has been used to extract the $\sin \phi$ Fourier component of the asymmetry.

² Note that in Ref. [8] a superscript on the asymmetry is used that is different and not related to the one here. Also the fit in Ref. [8] includes kinematic prefactors that are not needed here.

A possible uncertainty in the interpretation of the extracted asymmetries in terms of Eq. (2) is the contribution to the analyzed pion samples from the decay of exclusively produced vector mesons (VM). Due to the limited acceptance of the HERMES spectrometer, a large fraction of these vector mesons cannot be identified. Although the contribution of their decay pions to the observed pion yield is small – less than 15% for the highest z bin [8], based on a PYTHIA6 Monte Carlo simulation tuned for HERMES kinematics reproducing the exclusive VM cross section on a 10% level [34] – their contribution to $\langle \sin(\phi - \phi_S) \rangle_{UT}^q$ for a transversely polarized target could be significant [35]. For $\langle \sin \phi \rangle_{UL}^q$ this contributes only through the transverse component and is thus subtracted through Eq. (6). The VM contribution to the $\langle \sin \phi \rangle_{UL}^q$ moments from the longitudinal spin component of the target can be treated as a dilution as no $\sin \phi$ dependence on the longitudinal target polarization of either the VM production or its decay distribution is expected [36]. For an estimate of such effects moments were extracted that have the diluting contribution from this exclusive channel subtracted. This was done by dividing the $\langle \sin \phi \rangle_{UL}^q$ moments of Eq. (6) by $(1 - N_{VM}/N_{tot})$ where N_{VM} and N_{tot} are the numbers of pions from VM decays and all detected pions, respectively.

The main contribution to the systematic uncertainty in the extracted moments arises from the measurement of the target polarization. Other contributions include smearing due to detector resolution and radiative effects. The combined systematic uncertainty is found to be less than 0.003.

The moments for charged pions are shown as functions of x and z in Fig. 2 and summarized in Table I. In addition to the extracted longitudinal photon-axis moments $\langle \sin \phi \rangle_{UL}^q$ the lepton-axis moments for longitudinally and transversely polarized targets are plotted in Fig. 2. The latter include the prefactor $-\sin \theta_{\gamma^*}$ with which they appear in the $\langle \sin \phi \rangle_{UL}^1$ measurement. The resulting longitudinal photon-axis moments are significantly positive for the π^+ and consistent with zero for the π^- . Hence in the case of the π^+ this subleading-twist contribution dominates the measured lepton-axis asymmetries on a target that is polarized longitudinally with respect to the beam direction. Therefore it becomes clear that those asymmetries cannot be interpreted in terms of only the Collins fragmentation function or the Sivers function. In particular, the contribution from the Sivers function to the measured longitudinal lepton-axis asymmetries is small compared to the subleading-twist contribution as it appears only for the transverse component of the target spin. Unfortunately, due to the presence of several contributions (Eq. 2), it is not possible to make any statements about the size of any subleading-twist function separately. Nevertheless, it is clear that subleading-twist effects cannot be neglected a priori. This will be important when inter-




FIG. 2: The various azimuthal moments appearing in the measurement of the $\sin \phi$ modulations of single-spin asymmetries on a longitudinally polarized hydrogen target for charged pions as functions of x and z . The open symbols are the measured lepton-axis moments. The ones from a transversely polarized target are multiplied by $-\sin \theta_{\gamma^*}$ according to their appearance in the longitudinal lepton-axis moments. The closed symbol is the subleading-twist contribution to the measured lepton-axis asymmetries on a longitudinally polarized target. The triangles are slightly shifted horizontally for distinction. An overall systematic error of 0.003 is not included in the figure.

preting the measured lepton-axis asymmetries on a transversely polarized target which for experimental reasons receive not only contributions from the transverse target spin component (e.g., the Collins and Sivers effects) but also from the longitudinal component (subleading-twist) as in Eq. (1).³

In summary, single-spin asymmetries on hydrogen polarized longitudinally along the photon direction have been extracted for the first time. The contribution to the lepton-axis asymmetries from the transverse spin component in the measurement on a target polarized longitudinally with respect to the beam has been subtracted using the data from a transversely polarized hydrogen target. The averaged asymmetries in the range $0.023 < x < 0.4$ ($\langle x \rangle = 0.082$) and $0.2 < z < 0.7$ ($\langle z \rangle = 0.40$) are $0.030 \pm 0.004_{\text{stat}} \pm 0.002_{\text{sys}}$ for π^+ and $-0.009 \pm 0.006_{\text{stat}} \pm 0.001_{\text{sys}}$ for π^- . For π^+

³ It should be noted that a similar analysis for the azimuthal moments on a transversely polarized target yields corrections to the measured lepton-axis moments [8] that are negligible compared to their statistical uncertainty.

TABLE I: The $2\langle \sin \phi \rangle_{UL}^q$ moments of the π^+ and π^- production cross section for different x (top) and z (bottom) bins. Only statistical uncertainties are included. In addition there is a common systematic uncertainty of 0.003. Results are shown for all detected pions and for the case where the contribution from the decay of exclusive VM has been subtracted.

$\langle x \rangle$	$\langle z \rangle$	$\langle P_{h\perp} \rangle [\text{GeV}]$	$\langle y \rangle$	$\langle Q^2 \rangle [\text{GeV}^2]$	all π 's		π 's from exclusive VM subtracted	
					$2\langle \sin \phi \rangle_{UL}^{q,\pi^+}$	$2\langle \sin \phi \rangle_{UL}^{q,\pi^-}$	$2\langle \sin \phi \rangle_{UL}^{q,\pi^+}$	$2\langle \sin \phi \rangle_{UL}^{q,\pi^-}$
0.038	0.36	0.50	0.68	1.3	0.023 ± 0.008	-0.012 ± 0.010	0.025 ± 0.009	-0.013 ± 0.011
0.067	0.41	0.45	0.59	2.0	0.022 ± 0.007	-0.012 ± 0.010	0.023 ± 0.008	-0.012 ± 0.010
0.114	0.43	0.42	0.55	3.2	0.039 ± 0.010	-0.016 ± 0.013	0.041 ± 0.010	-0.017 ± 0.014
0.178	0.44	0.41	0.52	4.8	0.057 ± 0.016	0.028 ± 0.022	0.059 ± 0.016	0.029 ± 0.023
0.274	0.46	0.40	0.48	6.8	0.053 ± 0.025	-0.023 ± 0.035	0.054 ± 0.025	-0.024 ± 0.036
0.065	0.26	0.42	0.71	2.3	0.027 ± 0.009	0.000 ± 0.012	0.028 ± 0.009	0.000 ± 0.012
0.080	0.35	0.45	0.62	2.5	0.029 ± 0.008	-0.018 ± 0.011	0.030 ± 0.009	-0.019 ± 0.012
0.091	0.47	0.48	0.55	2.4	0.033 ± 0.008	-0.002 ± 0.011	0.035 ± 0.009	-0.002 ± 0.012
0.098	0.62	0.49	0.49	2.3	0.033 ± 0.011	-0.021 ± 0.015	0.037 ± 0.012	-0.024 ± 0.018

the $\langle \sin \phi \rangle_{UL}^q$ result is the dominating component in this range. This shows that subleading-twist effects are large and can, at HERMES kinematics, be comparable to leading-twist effects. This must be taken into account when interpreting asymmetries on transversely or longitudinally polarized targets solely in terms of leading-twist functions. At lower energies the isolation of leading-twist effects may be even more difficult.

We thank M. Diehl and F. Pijlman for many interesting discussions. We gratefully acknowledge the DESY management for its support and the staff at DESY and the collaborating institutions for their significant effort. This work was supported by the FWO-Flanders, Belgium; the Natural Sciences and Engineering Research Council of Canada; the National Natural Science Foundation of China; the Alexander von Humboldt Stiftung; the German Bundesministerium für Bildung und Forschung; the Deutsche Forschungsgemeinschaft (DFG); the Italian Istituto Nazionale di Fisica Nucleare (INFN); the Monbusho International Scientific Research Program, JSPS, and Toray Science Foundation of Japan; the Dutch Foundation for Fundamenteel Onderzoek der Materie (FOM); the U. K. Engineering and Physical Sciences Research Council and the Particle Physics and Astronomy Research Council; and the U. S. Department of Energy and the National Science Foundation.

- | | |
|--|---|
| <p>[1] J. P. Ralston, D. E. Soper, Nucl. Phys. B152 (1979) 109.</p> <p>[2] X. Artru, M. Mekhfi, Z. Phys. C 45 (1990) 669.</p> <p>[3] R. L. Jaffe, X. Ji, Nucl. Phys. B375 (1992) 527–560.</p> <p>[4] J. C. Collins, Nucl. Phys. B396 (1993) 161–182.</p> <p>[5] A. Airapetian, et al., Phys. Rev. Lett. 84 (2000) 4047–4051.</p> <p>[6] A. Airapetian, et al., Phys. Rev. D 64 (2001) 097101.</p> <p>[7] A. Airapetian, et al., Phys. Lett. B562 (2003) 182–192.</p> <p>[8] A. Airapetian, et al., Phys. Rev. Lett. 94 (2005) 012002.</p> <p>[9] V. Y. Alexakhin, et al., Phys. Rev. Lett. 94 (2005) 202002.</p> <p>[10] H. Avakian, et al., Phys. Rev. D69 (2004) 112004.</p> | <p>[11] E. Avetisyan, A. Rostomyan, A. Ivanilov, hep-ex/0408002.</p> <p>[12] D. W. Sivers, Phys. Rev. D 41 (1990) 83.</p> <p>[13] V. A. Korotkov, W. D. Nowak, K. A. Oganessyan, Eur. Phys. J. C18 (2001) 639–644.</p> <p>[14] M. Boglione, P. J. Mulders, Phys. Lett. B478 (2000) 114–120.</p> <p>[15] A. V. Efremov, K. Goeke, P. Schweitzer, Phys. Lett. B522 (2001) 37–48, erratum-ibid. B544 (2002) 389.</p> <p>[16] B.-Q. Ma, I. Schmidt, J. J. Yang, Phys. Rev. D66 (2002) 094001.</p> <p>[17] A. V. Efremov, K. Goeke, P. Schweitzer, Eur. Phys. J. C32 (2003) 337–346.</p> <p>[18] A. V. Efremov, K. Goeke, P. Schweitzer, Phys. Lett. B568 (2003) 63–72.</p> <p>[19] P. Schweitzer, A. Bacchetta, Nucl. Phys. A732 (2004) 106–124.</p> <p>[20] A. Bacchetta, U. D'Alesio, M. Diehl, C. A. Miller, Phys. Rev. D70 (2004) 117504.</p> <p>[21] M. Diehl, S. Sapeta, Eur. Phys. J. C41 (2005) 515–533.</p> <p>[22] A. Bacchetta, P. J. Mulders, F. Pijlman, Phys. Lett. B595 (2004) 309–317.</p> <p>[23] P. J. Mulders, R. D. Tangerman, Nucl. Phys. B461 (1996) 197–237, erratum-ibid. B484 (1996) 538.</p> <p>[24] D. Boer, P. J. Mulders, Phys. Rev. D57 (1998) 5780–5786.</p> <p>[25] K. Goeke, A. Metz, P. V. Pobylitsa, M. V. Polyakov, Phys. Lett. B567 (2003) 27–30.</p> <p>[26] H. Avakian, L. Elouadrhiri, AIP Conf. Proc. 698 (2004) 612–616.</p> <p>[27] S. J. Brodsky, D. S. Hwang, I. Schmidt, Phys. Lett. B530 (2002) 99–107.</p> <p>[28] J. C. Collins, Phys. Lett. B536 (2002) 43–48.</p> <p>[29] X. Ji, F. Yuan, Phys. Lett. B543 (2002) 66–72.</p> <p>[30] A. V. Belitsky, X. Ji, F. Yuan, Nucl. Phys. B656 (2003) 165–198.</p> <p>[31] A. Metz, M. Schlegel, Eur. Phys. J. A22 (2004) 489–494.</p> <p>[32] X. Ji, J.-P. Ma, F. Yuan, Phys. Rev. D71 (2005) 034005.</p> <p>[33] X. Ji, J.-P. Ma, F. Yuan, Phys. Lett. B597 (2004) 299–308.</p> <p>[34] P. Liebing, Desy-thesis-2004-036.</p> <p>[35] K. Goeke, M. V. Polyakov, M. Vanderhaeghen, Prog. Part. Nucl. Phys. 47 (2001) 401–515.</p> <p>[36] H. Fraas, Annals of Physics 87 (1974) 417–456.</p> |
|--|---|